



ISSN: 0975-833X

RESEARCH ARTICLE

MORPHO-ANATOMICAL CHARACTERISTICS OF LEAVES OF *TAXUS WALLICHIANA* ZUCC. FROM NORTH EASTERN AND NORTH WESTERN HIMALAYAS

^{*}, ¹Sharma, C. L., ¹Sharma, M., ²Singh, L., ¹Gogoi, B. R. and ²Kaur, M. J.

¹Department of Forestry, NERIST, Nirjuli-791109, Arunachal Pradesh, India

²Himalayan Research Group, Umesh Bhawan, Chotta Shimla, Shimla-171002, H. P., India

ARTICLE INFO

Article History:

Received 25th September, 2015
Received in revised form
08th October, 2015
Accepted 27th November, 2015
Published online 30th December, 2015

Key words:

Taxus wallichiana,
NE and NW Himalayas,
Stomatal density,
Spongy parenchyma, Palisade layer.

Copyright © 2015 Sharma et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Sharma, C. L., Sharma, M., Singh, L., Gogoi, B. R. and Kaur, M. J. 2015. "Morpho-anatomical characteristics of leaves of *Taxus wallichiana* Zucc. from North Eastern and North Western Himalayas", *International Journal of Current Research*, 7, (12), 24624-24627

INTRODUCTION

Taxus wallichiana Zucc. is a large slow growing evergreen tree of moderate height. It has great economic and medicinal values among gymnosperms and widely distributed in Northern Hemisphere. In India, it is found along the Himalayas between 2300m-3400m (Purkayastha, 1992). The bark and leaves of *T. wallichiana* are the prime source of taxol, a diterpene with exceptional anticancerous activity. It has also been reported that *Taxus wallichiana* also contain 10-deacytyle baccatin III, a precursor for the synthesis of taxol (Chattopadhyay et al., 1994). Since, a lot of work has been done on taxonomical, medicinal and conservational aspects of this species (Purohit et al., 1999, Khan et al., 2006, Spjut, 2007) but limited information is available on anatomical characteristics of leaves of *Taxus*. Spjut (2007) presented phytogeographic analysis of *Taxus* specimens on the basis of number of stomata rows in a stomata band and number of epidermal cells without papillae between leaf margin and stomatal band. It was debated over the years that *Taxus wallichiana* of NE states is better in quality and yield of taxol than the one found in NW Himalayas.

*Corresponding author: Sharma, C. L., Department of Forestry, NERIST, Nirjuli-791109, Arunachal Pradesh, India.

ABSTRACT

Leaves of *Taxus wallichiana* Zucc. from NE and NW Himalayas were studied to determine the variation in anatomical characteristics of both regions. The present study revealed that the leaves of NE Himalayas were larger in size than that of NW Himalayas. Anatomically, the spongy parenchyma of selected sites (except Morsing) of NE Himalaya did not contain pigmented cells. The number of palisade layers, stomatal bands and stomatal density were greater in leaves of NE Himalayas. Also, palisade cell size and vascular bundle size were larger in leaves of NE Himalayas as compared to NW Himalayas. Statistically, there were significant differences in palisade cell size, stomatal density and diameter of vascular bundle among selected sites of both regions.

Since, anatomical characters are well established criteria for identification of plants and herbal drugs. Therefore, the aim of present study is to study morpho-anatomical characteristics and variation in anatomical characteristics of leaves from both NE and NW Himalayas.

MATERIALS AND METHODS

Plant material

The leaves of *T. Wallichiana* Zucc. were collected from wild growing trees from NE and NW Himalayas. The selected sites along with geographical coordinates are listed in Table 1.

Table 1. Location of sites

Name of Site	Geographical Coordinates	Elevation
Shimla(NW Himalaya)	31 ^o 6' N, 77 ^o 10'E	2200m
Mandi (NW Himalaya)	31 ^o 43'N,76 ^o 55'E	2020m
Morsing (NE Himalaya)	27 ^o 08'N,92 ^o 13'E	2355m
Myodia (NE Himalaya)	28 ^o 14'N, 95 ^o 53'E	2360m
Bomdila (NE Himalaya)	27 ^o 16'N, 92 ^o 25'E	2234m

Determination of stomatal band and stomatal density

Fifteen fully mature leaves were selected from each site. The epidermis on abaxial side was separated from mesophyll by following the method of Locosselli and Ceccantini (2012). The permanent slides were prepared. Stomatal bands were determined by counting the rows of stomata between midrib and leaf margin. While, stomatal density was determined by using graph eye piece.

Morpho-anatomical Characteristics

For anatomical investigations, 25 fully mature leaves were selected from each site to measure their length and width. Some of leaves were preserved in 70% alcohol for anatomical investigations. The middle portion of the selected leaves were cut and processed for rotary microtomy. Sections of 8-10 μm thickness were cut and permanent slides were prepared by following standard laboratory protocol. A total of 50 leaves (10 leaves from each site) were taken for this parameter. Anatomical characteristics of leaves, palisade cell size and vascular bundle size were studied from the selected slides.

Table 2. Leaf anatomical characteristics of *Taxus wallichiana* from selected sites of North Eastern and North Western Himalayas

Parameter	Shimla	Mandi	Morsing	Myodia	Bomdila
Leaf length (mm) (Mean \pm SD)	27.27 \pm 1.89	26.28 \pm 1.77	32.81 \pm 2.32	33.16 \pm 3.18	29.95 \pm 3.69
Leaf width (mm) (Mean \pm SD)	2.37 \pm 0.19	2.19 \pm 0.22	2.81 \pm 0.26	3.4 \pm 0.16	2.55 \pm 0.13
Palisade layer (number)	1-2	1-2	1-2	1-3	1-3
Stomata band (number)	9-10	8-11	13-17	15-17	15-16
Stomata Density (Mean \pm SD)	30.80 \pm 1.99	31.60 \pm 2.55	39.00 \pm 2.94	37.50 \pm 3.14	31.30 \pm 2.11
Palisade cell size (μm) (Mean \pm SD)	34.84 \pm 5.99	33.93 \pm 7.04	58.50 \pm 10.14	68.77 \pm 15.07	66.82 \pm 12.86
Diameter of vascular bundle(μm) (Mean \pm SD)	244.20 \pm 12.67	256.83 \pm 9.41	425.25 \pm 13.10	309.46 \pm 10.12	393.67 \pm 6.86

Table 3. ANOVA of leaf anatomical characteristics of *Taxus wallichiana* from different sites of North Eastern and North Western Himalayas

Parameter		Sum of Squares	df	Mean Square	F value
Leaf length	Between Sites	983.227	4	245.807	34.292**
	Within Sites	860.172	120	7.168	
	Total	1843.399	124		
Leaf width	Between Sites	22.114	4	5.528	136.462**
	Within Sites	4.862	120	.041	
	Total	26.975	124		
Stomata density	Between Sites	605.320	4	151.330	22.654**
	Within Sites	300.600	45	6.680	
	Total	905.920	49		
Palisade cell size	Between Sites	23249.398	4	5812.349	50.008**
	Within Sites	11041.784	95	116.229	
	Total	34291.182	99		
Diameter of Vascular bundle	Between Sites	130890.761	4	32722.690	286.74**
	Within Sites	2282.412	20	114.121	
	Total	133173.173	24		

**= 1% at P < 0.01 level i.e. highly significant




Fig. 1. Stomatal bands between midrib and leaf margin

Statistical analysis

Statistical analysis was performed by using SPSS 18.0 software.

Photomicrography

Photomicrographs were taken with the help of image analysis system.

RESULTS AND DISCUSSION

Leaves play an important role in survival and growth of the plant. The leaves of both NE and NW Himalayas were needle

in shape. The leaves of NE Himalayas were larger in size than NW Himalayas (Table 2). According to Givnish (1984), the plants in the drier climate have smaller leaves to reduce evaporation and in more humid climates they have larger leaves because of available water. In the present study, the leaves of NE Himalayas were larger than NW Himalayas and is in agreement with the findings of Givnish (1984). Anatomically, *Taxus wallichiana* Zucc. leaves were covered by epidermis on both adaxial and abaxial surfaces. The epidermal cells were smaller and without papillose on adaxial surface whereas the epidermal cells were larger and papillose on the abaxial surface. Some of the leaves of NW Himalayas were revolute along the margin and no differentiation in the cell size of upper and lower epidermal was observed. The stomata were hypostomatic. Below epidermis, mesophyll was differentiated into palisade and spongy parenchyma. Palisade parenchyma cells were elongated and arranged in 1- 3 rows beneath adaxial




Fig. 2. T. S. of leaves collected from different sites of North East and North West Himalayas

epidermis. The spongy parenchyma cells were more or less oval in shape and loosely arranged. There was only one having xylem on the upper side and phloem on lower side. A few cells of transfusion tissue were also present on the side of xylem and phloem. The anatomical study of leaves is in agreement with the previous anatomical studies (Hamidipour *et al.*, 2011). In the present study, the parenchyma cells were filled with dark red or yellowish brown deposits in both epidermal layers, vascular bundle sheath and cells of spongy parenchyma in the leaves of both sites of NW Himalayas and in one site (Morsing) of NE Himalayas (Fig. 2.). While, these pigmented cells were found absent in vascular bundle sheath and spongy parenchyma of other selected sites of NE Himalayas.

The stomata were arranged in 8-11 rows between leaf margin and mid rib in leaves of NW Himalayas while there were 13-17 rows in leaves of NE Himalayas (Fig. 1). The available literature reveals that plants growing in more stressful environment conditions have more stomatal density (Dinis *et al.*, 2011). On contrary to this, Locosseli and Ceccantini (2012) report higher stomatal density and stomatal index in *Podocarpus lambertii* growing in warmer and drier climates than individuals growing in colder and rainier site. The results given in Table 2 shows that stomatal density in selected sites of NW Himalayas were more or less same. On the other hand, the maximum stomatal density was observed in Morsing (39.00 ± 2.94) among selected sites of NE Himalayas. There was significant difference in stomatal density of both region (Table 3). The difference in stomatal density may be due to larger size of leaves of NE Himalayas. There was not much variation in number of palisade layers of both regions. However, the palisade cell size varied from $33.93 \pm 7.04 \mu\text{m}$ to $34.84 \pm 5.99 \mu\text{m}$ in selected sites of NW Himalayas and from $58.50 \pm 10.14 \mu\text{m}$ to $68.77 \pm 15.07 \mu\text{m}$ in NE Himalayas (Table 2). Likewise, the size of vascular bundles varied from $244.20 \pm 12.67 \mu\text{m}$ to $256.83 \pm 9.41 \mu\text{m}$ in NW Himalayas and $309.46 \pm 10.12 \mu\text{m}$ to $425.13.10 \mu\text{m}$ in NE Himalayas (Table 2). There was significant differences in palisade cell size and vascular bundle size of both the regions (Table 3). The significant differences in these parameters may be due to ecological adaptations of the plant to different climatic conditions.

Conclusions

The present study revealed morphologically the leaves are of same shape and different sizes. Anatomically, the leaves of NE

and NW Himalayas have uniform structure. The number of palisade layer, stomatal bands, stomatal density, size of palisade cells and vascular bundles are greater in NE Himalayas than NW Himalayas, which might be due to differences in ecological conditions.

Acknowledgements

The authors are grateful to Department of Biotechnology, New Delhi for financial assistance.

REFERENCES

- Chattopadhyay, S.K., Tripathi, V.K., Thakur, R.S., Sharma R.P. and Jain S.P. 1994. Isolation of taxol, 10-deacetyl baccatin III and (-) betuligenol from *Taxus baccata*. *Ind. J. Chem.*, 33B: 409–411.
- Dinis, L.T., Peixoto, F., Pinto, T., Costa, R., Bennet, R. N. And Gomes-Laranjo, J. 2011. Study of morphological and phenological diversity in chestnut trees ('Judia' variety) as a function of temperature sum. *Env. Exp. Botany*, 70:110-120.
- Givnish, T. J. 1984. Leaf and canopy adaptations in tropical forests. In Medina E., Mooney H. A. , Vazquez-Yanés C., eds. Physiological ecology of plants of wet tropics. The Hague, the Netherlands: Dr. Junk, 51-84.
- Hamidipour, A., Radjabian, T., Charlet D.A. and Zarrei M. 2011. Leaf anatomical investigation of Cupressaceae and Taxaceae in Iran. *Wulfenia*, 18: 95 –111.
- Khan, M., Verma, S.C., Srivastava, S.K., Shawl, A.S., Syamsundar, K.V., Khanuja, S.P.S. and Kumar, T. 2006. Essential oil composition of *Taxus wallichiana* Zucc. from the Northern Himalayan region of India. *J. Flavour Frag.*, 21:772-775.
- Locosselli, G. M. And Ceccantini, G. 2012. Plasticity of stomatal distribution pattern and stem tracheid dimensions in *Podocarpus lambertii*: an ecological study. *Annals of Bot.*, 110:1057-1066.
- Purkayastha, S. K. 1992. A manual of Indian timbers, Sribhumi Publications, Calcutta.
- Purohit, A., Maikhuri, R.K., Rao, K.S. and Nautiyal, S. 2001. Impact of bark removal on survival of *Taxus baccata* L. (Himalayan yew) in Nanda Devi Biosphere Reserve, Garwhal Himalaya, India. *Curr. Sci.*, 81:586–590.
- Spjut R.W. 2007. A phytogeographical analysis of *Taxus* (Taxaceae) based on leaf anatomical characters. *J. Bot. Res., Inst. Texas*, 1: 291 – 332.
